

# When she talks to you about the violence

**A toolkit  
for GPs  
in NSW**

**Over 1 in 5 women make their first disclosure of domestic violence to their GP.<sup>1</sup>**

**You may be the only person she will tell.**

**Your skills and sensitivity are essential.**

This resource has been developed to assist you in identifying and responding to women and children who have experienced or are experiencing family violence (also known as 'domestic violence' or 'intimate partner violence')

*'It has been estimated that full time GPs are seeing up to five women per week who have experienced some form of intimate partner abuse (physical, emotional or sexual) in the past 12 months.'<sup>2</sup>*

The toolkit contains guidelines for patient care, from a range of sources, as well as some legal information relevant to your role as her GP.

*'The Medical Profession has key roles to play in early detection, intervention and provision of specialized treatment of those who suffer the consequences of domestic violence, whether it be physical, sexual or emotional.'<sup>3</sup>*

Responding effectively to family violence requires knowledge of the physical and emotional consequences of the violence, an understanding of appropriate and inappropriate responses, and having good networks with local family violence services.

*After family and friends, victims are most likely to tell health professionals about violence.<sup>4</sup>*

## Contents

1. What is family violence?
2. Indicators
3. How to ask your patient
4. Responding to a disclosure
5. Initial safety planning
6. Victims support scheme
7. Note-taking for legal purposes
8. Mandatory reporting
9. Immigration family violence provisions
10. Continuing care
11. When your patient is the perpetrator
12. When both partners are your patients
13. Subpoenas
14. Referrals
15. Training & resources



This toolkit has been developed by Women's Legal Services NSW, with the generous assistance of several individuals and organisations. It updates and extends 'Identifying Family Violence: A Resource Kit for General Practitioners in New South Wales,' 1999, Domestic Violence Advocacy Service NSW.

Information about the law is presented in summary form and should not be relied upon as a substitute for professional legal advice.

Organisations have permission to reproduce parts or the whole of the publication for the purposes of workshops, seminars, etc as long as the original meaning is retained and proper credit given.



This publication has been produced with the financial assistance of the Law and Justice Foundation of NSW. The Foundation seeks to advance the fairness and equity of the justice system and to improve access to justice, especially for socially and economically disadvantaged people. [www.lawfoundation.net.au](http://www.lawfoundation.net.au)

Disclaimer: any opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the Law and Justice Foundation's Board of Governors.

Published in October 2013.

Available online at <http://itstimetotalk.net.au/gp-toolkit>

## 1. What is family violence?

Family or domestic violence is **an abuse of power** within a close relationship, or after separation. It involves one person dominating and controlling another, causing **intimidation and fear**.

It is **not necessarily physical** and can include:

- sexual abuse,
- emotional or psychological abuse,
- verbal abuse,
- spiritual abuse,
- stalking and intimidation,
- social and geographic isolation,
- financial abuse,
- cruelty to pets, or
- damage to property.

Often the terms 'family violence' and 'domestic violence' are used interchangeably. 'Family violence' is sometimes thought of as the broader term, covering intimate, family and other relationships of mutual obligation and support.

Family violence is **often experienced as a pattern of abuse** that escalates over time.

Most domestic violence is perpetrated by men, against women and children.<sup>5</sup> However women can also be perpetrators of violence, and domestic violence also happens in same-sex relationships.

Women are at greater risk of violence from intimate partners during pregnancy, or after separation. A safety survey conducted by the Australian Bureau of Statistics in 2005 found that 17% of women who had experienced violence from a partner during a relationship, experienced it for the first time during pregnancy.

## 2. Indicators

*'When assessing your patient... remember that most presentations of family violence are probably hidden and not the obvious black eye.'*<sup>6</sup>

The following are indicators associated with victims of family violence.

Indicators in adults	Indicators in children
<p><b>Physical</b></p> <ul style="list-style-type: none"> <li>• Unexplained bruising and other injuries</li> <li>• Bruises of various ages</li> <li>• Head, neck and facial injuries</li> <li>• Injuries on parts of the body hidden from view (including breasts, abdomen and/or genitals), especially if pregnant</li> <li>• 'Accidents' occurring during pregnancy</li> <li>• Miscarriages and other pregnancy complications</li> <li>• Injuries to bone or soft tissues</li> <li>• Injuries sustained do not fit the history given</li> <li>• Bite marks, unusual burns</li> <li>• Chronic conditions including headaches, pain and aches in muscles, joints and back</li> <li>• Ulcers</li> <li>• Dizziness</li> <li>• Sexually transmitted disease</li> <li>• Other gynaecological problems</li> </ul> <p><b>Psychological/behavioural</b></p> <ul style="list-style-type: none"> <li>• Emotional distress, eg, anxiety, indecisiveness, confusion, and hostility</li> <li>• Sleeping and eating disorders</li> <li>• Anxiety/depression/pre-natal depression</li> <li>• Psychosomatic and emotional complaints</li> <li>• Drug abuse</li> <li>• Self-harm or suicide attempts</li> <li>• Evasive or ashamed about injuries</li> <li>• Multiple presentations at the surgery/client appears after hours</li> <li>• Partner does most of the talking and insists on remaining with the patient.</li> <li>• Seeming anxious in the presence of the partner.</li> <li>• Reluctance to follow advice</li> <li>• Social isolation/no access to transport</li> <li>• Frequent absences from work or studies</li> <li>• Submissive behaviour/low self esteem</li> <li>• Alcohol or drug abuse</li> </ul>	<p><b>Physical</b></p> <ul style="list-style-type: none"> <li>• Difficulty eating / sleeping</li> <li>• Slow weight gain (in infants)</li> <li>• Physical complaints</li> <li>• Eating disorders</li> </ul> <p><b>Psychological / behavioural</b></p> <ul style="list-style-type: none"> <li>• Aggressive behaviour and language</li> <li>• Depression, anxiety and/or suicide attempts</li> <li>• Appearing nervous and withdrawn</li> <li>• Difficulty adjusting to change</li> <li>• Regressive behaviour in toddlers</li> <li>• Delays or problems with language development</li> <li>• Psychosomatic illness</li> <li>• Restlessness and problems with concentration</li> <li>• Dependent, sad or secretive behaviours</li> <li>• Bedwetting</li> <li>• 'Acting out', for example cruelty to animals</li> <li>• Noticeable decline in school performance</li> <li>• Fighting with peers</li> <li>• Over protective or afraid to leave mother</li> <li>• Stealing and social isolation</li> <li>• Abuse of siblings or parents</li> <li>• Alcohol and other drug use</li> <li>• Psychosomatic and emotional complaints</li> <li>• Exhibiting sexually abusive behaviour</li> <li>• Feelings of worthlessness</li> <li>• Transience</li> </ul>

Figure 1: Indicators associated with victims of family violence.

*'Frequently there are no visible signs of assault or rape in domestic violence presentations. This does not mean that the emotional or psychological effects of the assault are... any less devastating to the victim.'*<sup>7</sup>

### 3. How to ask your patient

*'In any situation that you suspect underlying psychosocial problems you can ask indirectly and then directly about partner abuse.'*<sup>8</sup>

If you have concerns that your patient is experiencing family violence, you should ask to speak with her alone, separate from her partner or any other family members.

You can always ask **broad questions** about whether your patient's relationships are affecting her health and wellbeing. For example:

- 'How are things at home?'
- 'How are you and your partner getting on?'
- 'Is anything else happening which might be affecting your health?'

*'It is important to realise that women who have been abused want to be asked about domestic violence and are more likely to disclose if asked.'*<sup>9</sup>

If appropriate, you can ask **direct questions** about any violence. For example:<sup>10</sup>

- 'Are there ever times when you are frightened of your partner?'
- 'Are you concerned about your safety or the safety of your children?'
- 'Does the way your partner treats you make you feel unhappy or depressed?'
- 'Has your partner ever physically threatened or hurt you?'
- 'Violence is very common in the home. I ask a lot of my patients about abuse because no one should have to live in fear of their partners.'

If you see **specific clinical symptoms**, you can ask **specific questions** about these (eg, bruising). These could include:

- 'You seem very anxious and nervous. Is everything alright at home?'
- 'When I see injuries like this, I wonder if someone could have hurt you?'
- 'Is there anything else that we haven't talked about that might be contributing to this condition?'

If your patient's fluency in English is a barrier to discussing these issues, you should work with a qualified interpreter. Don't use her partner, other family members or a child as an interpreter. It could compromise her safety, or make her uncomfortable to talk with you about her situation. The Doctors' Priority Line, phone 1300 575 847, is a 24/7 free telephone interpreting service to assist GPs to communicate with patients from non-English speaking backgrounds.

### 4. Responding to a disclosure

Your immediate response and attitude when a woman discloses family violence can make a difference.

*'Patients... value emotional support from healthcare professionals, careful and non-judgmental listening, and reassurance that the abuse is not their fault and that negative feelings are understandable.'*<sup>11</sup>

#### Listen

Being listened to can be an empowering experience for a woman who has been abused.

#### Communicate belief

'That must have been frightening for you.'

#### Validate the decision to disclose

'I understand it could be very difficult for you to talk about this.'

#### Emphasise the unacceptability of violence

'Violence is unacceptable; you do not deserve to be treated this way.'

#### Be clear that she is not to blame

Avoid suggesting that the woman is responsible for the violence or that she is able to control the violence by changing her behaviour.

#### Do not ask

- 'Why don't you leave?'
- 'What could you have done to avoid this situation?'
- 'Why did he hit you?'



Figure 2: Key steps after a disclosure of family violence

## 5. Initial safety planning

Assist your patient to evaluate her immediate and future safety, and that of her children. Best-practice risk assessment involves seeking relevant facts about her particular situation, asking her about her own perception of risk, and using professional judgment. You may need to refer your patient to a specialised domestic violence service such as the Domestic Violence Line. See 'Abuse and violence: Working with our patients in general practice' (white book) for detailed guidance on your role as a GP.




Figure 3: Aspects of best-practice risk assessment

For initial safety planning, you will at least need to:

- Speak to the woman alone
- Check for **immediate concerns**
  - Does she feel safe going home after the appointment?
  - Are her children safe?
  - Does she need an immediate place of safety?
  - Does she need to consider an alternative exit from your building?
- If immediate safety is not an issue, check her **future safety**
  - Does he have weapons?
  - Does she need a referral to police or a legal service to apply for an Apprehended Violence Order?
  - Does she have **emergency telephone numbers**?
    - NSW Police: 000 or 106 (TTY)
    - Domestic Violence Line (24/7 emergency, referral and counselling line for people experiencing

domestic violence. Can explain basic information about AVOs and assist with risk assessment): 1800 656 463 or 1800 671 442 (TTY)

- Does she need a referral to a domestic violence service to help make **an emergency plan**:  
Where would she go if she had to leave?  
How would she get there?  
What would she take with her?  
Who are the people she could contact for support?
- Document any plans made, for future reference.

Risk assessment is an ongoing process. You may need to check in on your patient to follow up on this initial safety plan. See section 9 (Continuing care).

*'It is important to remember that the true goal... is to prevent violence, not predict it.'*<sup>1,2</sup>

## 6. Victims support scheme

Victims of violent crime that occurred in NSW may be entitled to various forms of support through the victims support scheme, administered by Victims Services NSW.

If your patient has experienced family violence, she can apply for free counselling. She may also be eligible for financial assistance for her immediate needs (such as relocation expenses, if required for her safety, or emergency medical and dental expenses.) Further financial assistance for economic loss can be applied for to cover ongoing costs such as medical or dental expenses, loss of earnings. Financial assistance is capped to set limits. Some victims may also be granted a recognition payment depending on the nature of violence that occurred.

Most types of support require victims to apply within set time frames (usually within 2 years from the incident).

Your patient may need you to write a medical report or provide evidence of injuries she suffered, if she is applying for financial assistance or a recognition payment. Applications for free counselling only do not need any supporting documentation.

Refer your patient to Victims Services NSW for more information. A support co-ordinator will assist her to apply for any benefits available under this scheme.

## 7. Note-taking for legal purposes

Your notes may be required as evidence, if charges are laid against the perpetrator.

If family violence is a concern, you should keep detailed notes that:

- **Describe physical injuries** (including the type, extent, age and location). If you suspect violence is a cause, but your patient has not confirmed this, include your comment as to whether her explanation accurately explains the injury.
- **Record what the patient said** (using quotation marks)
- **Record any relevant behaviour observed**, being detailed and factual rather than stating a general opinion, eg, rather than 'the patient was distressed', write 'the patient cried throughout the appointment, shook visibly and had to stop several times to collect herself before answering a question'.

**Consider taking photographs of injuries**, or certifying photographs taken of the injuries presented at the time of consultation.

To be good evidence in court, file notes must include date and time, and clearly identify the client. You must clearly identify yourself as the author, and sign the file note. Do not include generalisations or unsubstantiated opinions. Correct and initial any errors, set out your report sequentially, and use only approved symbols and abbreviations.

## 8. Mandatory reporting

If a patient talks about experiencing or perpetrating violence, you may need to report this to Community Services. You have an obligation to report if you believe you have reasonable grounds to suspect that a child is at risk of significant harm.

Exposing children to domestic violence can have a serious psychological impact on children. In some cases you may feel there is risk of significant harm to a child even though it seems unlikely that the violent person in their home would physically hurt them. Use your professional judgment about the individual circumstances and the nature of the violence.

Community Services has online resources for mandatory reporters at [www.community.nsw.gov.au/docs\\_menu/preventing\\_child\\_abuse\\_and\\_neglect/resources\\_for\\_mandatory\\_reporters.html](http://www.community.nsw.gov.au/docs_menu/preventing_child_abuse_and_neglect/resources_for_mandatory_reporters.html). These include a Mandatory Reporting Guide to assist GPs and other mandatory reporters with making 'risk of significant harm' assessments.

## 9. Immigration family violence provisions

There are special family violence provisions in immigration law that are intended to relieve the fear of a 'partner visa' applicant who may believe that she needs to stay in an abusive relationship in order to remain in Australia. These provisions allow certain applicants to obtain permanent residence even if the relationship with their Australian sponsor has broken down, where there is evidence of family violence against the applicant or her dependent child.

A report or statutory declaration from a GP detailing physical injuries and/or treatment for mental health issues that are consistent with family violence can be used as part of the evidence given to the Department of Immigration and Citizenship to access the provisions.

If your patient has concerns about her visa to stay in Australia, you may wish to refer her to Immigrant Women's Speakout, phone (02) 9635 8022, or the Immigration Advice & Rights Centre, phone (02) 9279 4300.

## 10. Continuing care

- Consider your patient's safety as a paramount issue. A woman is usually a good judge of her own safety. You can help to monitor the safety of her and her children by asking about any escalation of violence.
- Empower her to take control of decision-making; ask what she needs and present choices of actions she may take and services available.
- Respect the knowledge and coping skills she has developed. You can help build on her emotional strengths, for example, by asking 'How have you dealt with this situation before?'
- Provide emotional support.
- Ensure confidentiality – the woman may suffer additional abuse if her partner suspects she has disclosed the abuse.
- Be familiar with appropriate referral services and their processes. Patients may need your help to seek assistance. Have information available for the patient to take with her if appropriate.

*'I dropped some hints to test the water. [The GP] was supportive without being interfering and because of this I made the decision to tell her. She was fantastic and told me about the [Domestic Violence Line] who I called and put me into contact with a women's refuge. I am rebuilding my life, and looking forward to a happy future.'*<sup>13</sup>

## 11. When your patient is the perpetrator

Consider the safety of female victims and their children as the highest priority. Note that perpetrators of violence have a tendency to minimise the violence, or shift blame.

If violence is suspected and further information is needed, start with broad questions such as:

- 'How are things at home?'

Then if violence is disclosed, ask more specific questions such as:

- 'Some men who are stressed like you hurt the people they love. Is this how you are feeling? Did you know that there are services that can help you?'

Acknowledge the existence of violence by statements such as:

- 'That was brave of you to tell me. Sometimes people who are stressed hurt the people they love. However, violent behaviour towards your partner and other family members is never acceptable. It not only affects your partner but your children as well. Did you know there are services which may be able to assist you?'

## 12. When both partners are your patients

Special care is required if a patient discloses family violence, and the violent person is also your patient or is a patient within the same service.

If you have seen the victim or her children, your primary duty is to them. If the perpetrator is also your patient, you should refer them to another practitioner or another practice.

If both partners remain within your practice, you will need to take extra caution, for example:<sup>14</sup>

- Establish staff protocols that ensure confidentiality of records.
- There should be no discussion about suspected or confirmed abuse with the violent partner unless the woman consents to it.
- If a woman agrees that you can talk with her partner about the violence, it is important that a safety plan is in place.

**Couple or marital counselling is not appropriate** in circumstances where there has been domestic violence, due to the power imbalance in the relationship and the threat to the woman's safety.

## 13. Subpoenas

As a GP, you could be served with a subpoena relating to a patient. Where family violence is present, dealing with a subpoena requires even more care than usual. For example, a woman who has experienced sexual assault may be able to claim the *Sexual Assault Communications Privilege* to maintain the confidentiality of your records if subpoenaed.

A subpoena is a stamped court order to hand over documents (a subpoena to produce), to attend court as a witness (a subpoena to give evidence) or both (a subpoena to produce and give evidence). Subpoenas are issued as part of a court case such as a criminal law proceeding or a family law dispute, at the request of one of the parties.

It is important to treat subpoenas with caution, especially when the person seeking the information is not your patient, eg, is her ex-partner. First, check that the subpoena is valid: has a court stamp, has been served on you before the stated deadline and that conduct money has been provided.

You must *respond* to a valid subpoena – either to obey the orders, or to object. There are various grounds for objecting to a subpoena, for example: the request is too onerous, or the information is 'privileged' (protected by law).

*Always contact your patient* to let her know that you have been served with a subpoena, and to ask her how she would like you to respond. Note that you may be legally required to go against her wishes.

Subpoenas requesting documents will have a schedule of what material is being sought. *Never hand over more* than what is listed in this schedule.

In some cases, you or your patient may need legal advice. You could seek guidance from the AMA, the RACGP, your insurer, or a private lawyer. Your patient could get legal advice from her own lawyer, a community legal centre, or – if appropriate – the Sexual Assault Communication Privilege Service at Legal Aid.

## 14. Referrals

Here are some key contacts for patients.

<b>Domestic Violence Line</b>	1800 656 463 or 1800 671 442 (TTY) <a href="http://www.community.nsw.gov.au">www.community.nsw.gov.au</a>	24/7 emergency, referral and counselling line for people experiencing domestic violence. Can explain basic information about AVOs and assist with risk assessment.
<b>MensLine Australia</b>	1300 78 99 78 <a href="http://www.mensline.org.au">www.mensline.org.au</a>	24/7 support, information and referral service, helping men deal with relationship problems.
<b>Law Access</b>	1300 888 529 <a href="http://www.lawaccess.nsw.gov.au">www.lawaccess.nsw.gov.au</a>	Free legal information, referrals and in some cases, advice for people who have a legal problem in NSW
<b>Victims Services</b>	1800 633 063 <a href="http://www.lawlink.nsw.gov.au/vs">www.lawlink.nsw.gov.au/vs</a>	Support, referrals and information for victims of crime in NSW. Support coordinators help victims apply for assistance under the NSW victims support scheme.
<b>Legal Aid NSW – Sexual Assault Communication Privilege Service</b>	<a href="mailto:sacps@legalaid.nsw.gov.au">sacps@legalaid.nsw.gov.au</a> <a href="http://www.legalaid.nsw.gov.au/what-we-do/civil-law/sexual-assault-communications-privilege-service">http://www.legalaid.nsw.gov.au/what-we-do/civil-law/sexual-assault-communications-privilege-service</a>	Free legal advice for victims of sexual assault if their records are the subject of a subpoena in criminal proceedings.

See <http://itstimetotalk.net.au/gp-toolkit> for more extensive referrals.

## 15. Training & resources

- **RACGP – Abuse and violence: Working with our patients in general practice (white book)**  
[www.racgp.org.au/your-practice/guidelines/abuse-and-violence](http://www.racgp.org.au/your-practice/guidelines/abuse-and-violence)
- **AFP – Intimate partner violence: Identification and response in general practice**  
[www.racgp.org.au/afp/2011/november/intimate-partner-violence](http://www.racgp.org.au/afp/2011/november/intimate-partner-violence)
- **NSW Health – The Education Centre Against Violence**  
[www.ecav.health.nsw.gov.au](http://www.ecav.health.nsw.gov.au)

## References

1. Jo Spangaro & Anthony Zwi, *After the Questions: Impact of Routine Screening for Domestic Violence in NSW Health Services*, (School of Public Health and Community Medicine, The University of New South Wales, 18 August 2010), 22.
2. Gwenneth Roberts et al, *Intimate Partner Abuse and Health Professionals: New Approaches to Domestic Violence*, (London: Churchill Livingstone Elsevier, 2006), 19–40, cited in Jane London, *Abuse and Violence: Working with Our Patients in General Practice* (The Royal Australian College of General Practitioners, 3rd ed, 2008) 11.
3. Australian Medical Association, *AMA Position Statement on Domestic Violence*, (Canberra, AMA, 1998) 1.
4. Keys Young, *Against the Odds: How Women Survive Domestic Violence*, (Office of the Status of Women, 1998).
5. Access Economics & Commonwealth of Australia, *The Cost of Domestic Violence to the Australian Economy: Part II*, (Office for the Status of Women, 2004) 9.
6. Kelsey Hegarty, 'Identification and management role of GPs for women experiencing partner abuse' (Paper presented at the 25th Congress Medical Women's International Association, Sydney, 19 April 2001) 1, available at <[http://www.regional.org.au/au/mwia/papers/full/17\\_hegarty.htm](http://www.regional.org.au/au/mwia/papers/full/17_hegarty.htm)>
7. Jane London, above n 2, at 7.
8. Kelsey Hegarty, above n 6, at 1.
9. Kelsey Hegarty, 'The hidden epidemic of domestic violence: when to ask and how to respond' (2012) 13(7) *Medicine Today* 54–57; Kelsey Hegarty & Lorna O'Doherty, 'Intimate partner violence – identification and response in general practice' (2011) 40(11) *Aust Fam Physician* 852–6.
10. Kelsey Hegarty, above n 6, at 1.
11. Charles George, *Domestic Violence: A Report from the BMA Board of Science*, (British Medical Association, 2007) 35.
12. Don G Dutton & P Randall Kropp, 'A review of domestic violence risk instruments', (2000) 1(2) *Trauma, Violence and Abuse*, 171–181 at 179.
13. Charles George above n 10 at 36.
14. Based on Lorraine E. Ferris et al., 'Guidelines for managing domestic abuse when male and female partners are patients of the same physician', (1997) 278 (10) *Journal of the American Medical Association*, 851–857.